

Experimental Particle Physics PHYS6011

Looking for Higgs and SUSY at the LHC

or...what can you get for \$10,000,000,000

Lecture 5

**First LHC collisions at 3.5 TeV
per beam 30th March 2010**

ATLAS
EXPERIMENT
<http://atlas.ch>

Search for the Higgs Boson

- Missing piece of Standard Model
- Standard Model Higgs theory well understood:
 - Mass is only free parameter
 - Clear predictions to test
- Most “New Physics” models have something equivalent to a Higgs boson (“MSSM Higgs”, “little Higgs”, etc...).
- Could be more than one type of Higgs boson
- Current limit $M_H > 115$ GeV (LEP)
- Particle masses are generated by interactions with the scalar (Higgs) field.
- Couplings are fixed by the masses.
- Once M_H is known everything is predicted.
- So by measuring the coupling of the Higgs to particles of known mass we can test theory.


Higgs Mechanism in the Standard Model

- Need to accommodate massive gauge bosons

- Strong and electromagnetism ok (photon, gluon)
- Weak force has a massive W and Z

Modified potential $V = \mu^2 |\phi|^2 + \lambda |\phi|^4$

$$v = \sqrt{\frac{-\mu^2}{\lambda}}$$



- **Step 1: Spontaneous Symmetry Breaking** produces one massive and one massless gauge boson (**Goldstone Boson**).
- Step 2: Introduce **local gauge invariance** : massive Higgs particle and a massive gauge field.

- Higgs mass a free parameter

$$M_H = \sqrt{-2\mu^2}$$

- Gauge couplings of Higgs doublet give gauge boson masses



$$M_W = g_W v / 2 \quad M_Z = M_W \cos \theta_W \quad \cos \theta_W = 0.8810$$

- Higgs couplings to fermions depends on their mass and unique coupling for each fermion:

$$M_f \propto M_H g_f$$

What do we know about the Higgs?


- No useful lower limit from theory.
- Upper limit from WW scattering
 - Above $\sim 1\text{TeV}$ cross-section $\rightarrow \infty$
 - Need Higgs to “regularise” cross-section



What do we know about the Higgs?

If no new physics up to Planck scale ($\sim 10^{19}\text{GeV}$)
small mass range for Higgs: $130 < M_H < 190 \text{ GeV}$

$$M_H^2 \leq \frac{8\pi^2 v^2}{3 \log \frac{\Lambda_{QCD}^2}{v^2}} \quad v^2 = 246 \text{ GeV}$$




What do we know about the Higgs?

Direct searches
 $M_H > 114.4 \text{ GeV}$
 @ 95% C.L.

Search for the Higgs Particle

Status as of March 2009


Excluded by
 LEP Experiments
 95% confidence level



90% confidence
 95% confidence

Excluded by
 Tevatron
 Experiments

Excluded by
 Indirect Measurements
 95% confidence level



<http://lepewwg.web.cern.ch/LEPEWWG>

How to discover a signal?





- Total number of events (n_t) will have signal events (n_s) and background events (n_b)
- Number of events follows a Poissonian distribution with $\sigma = \sqrt{n}$.
- Require signal $> 5\sigma$ above background for “**observation**”.

$$\text{Significance } S = n_s / \sqrt{n_b} > 5$$

- Require signal $> 3\sigma$ above background for “**first evidence**”.
- e.g. Measure 140 events and know 100 come from background:
$$S = 40 / \sqrt{100} = 4$$
- How do you know the background? Monte Carlo or Look in areas where there is no signal.
- Significance depend on how much data you have taken

$$S \propto \sqrt{\text{Luminosity}}$$


Higgs Production Mechanisms




Gluon fusion most promising

Higgs production and decay

How often is it produced?





What does the Higgs decay into?



Process	Events/s	Events/year
$W \rightarrow e\nu$	40	$4 \cdot 10^8$
$Z \rightarrow ee$	4	$4 \cdot 10^7$
$t\bar{t}$	1.6	$1.6 \cdot 10^7$
$b\bar{b}$	10^6	10^{13}
$\tilde{g}\tilde{g}$ ($m = 1 \text{ TeV}$)	0.002	$2 \cdot 10^4$
Higgs ($m = 120 \text{ GeV}$)	0.08	$8 \cdot 10^5$
Higgs ($m = 120 \text{ GeV}$)	0.08	$8 \cdot 10^5$
Higgs ($m = 800 \text{ GeV}$)	0.001	10^4
QCD jets $p_T > 200 \text{ GeV}$	10^2	10^9

Higgs Decay

Detectable decays of a Higgs-Boson




Branching Fraction: If produce 10^8 Higgs and measure only 20 decays $H \rightarrow gg$ with an efficiency of 0.00025% then Branching Fraction:

$$BF(Higgs \rightarrow gg) = \frac{N_{decays}}{N_H * \eta} = \frac{20}{10^8 * 2.5 \times 10^{-6}} = 0.08$$


- Which decay to look at?
- Depends on Higgs Mass
 - $M_H < 100 \text{ GeV}$
 - $gg \rightarrow H \rightarrow \gamma\gamma$
 - $M_H < 150 \text{ GeV}$
 - $H \rightarrow ZZ^* \rightarrow 4l$
 - $H \rightarrow bb$
 - $H \rightarrow WW^* \rightarrow 2l 2\nu$
 - $H \rightarrow \tau^+\tau^-$
 - $130 < M_H < 500 \text{ GeV}$
 - $H \rightarrow ZZ \rightarrow 4l$
 - $600 < M_H < 1000 \text{ GeV}$
 - $H \rightarrow ZZ, WW \rightarrow \text{jets}$

Best Modes to look at



Compare to list on previous slides

Backgrounds - Tevatron to the LHC



Huge stats for Standard Model signals. Rates @ $10^{33} \text{ cm}^{-2} \text{ s}^{-1}$

$\sim 10^9$ events/ 10 fb^{-1} W (200 Hz)

$\sim 10^8$ events/ 10 fb^{-1} Z (50 Hz)

$\sim 10^7$ events/ 10 fb^{-1} tt (1 Hz)

($10 \text{ fb}^{-1} = 1$ year of LHC running at low luminosity $10^{33} \text{ cm}^{-2} \text{ s}^{-1}$)

Background is anything with signature similar to signal


- $W+X$ (X can be W , Z or just 2 QCD jets)
- $ZZ \rightarrow qql+l-$ (one lepton not identified)
- $\tau^+\tau^-$
- b -tags can be real, charm or fakes

Current Results - Tevatron


- At 120 GeV $H \rightarrow b\bar{b}$ dominates
- Signature $gg \rightarrow H \rightarrow b\bar{b}$:
 - 2 jets
 - One or two b -tags
- Swamped by dijet production
 - $b\bar{b} \sim \mu\text{b}$
 - $q\bar{q} \sim \text{mb}$ (*fake b -tag rate small but not zero*)
- Have to use $W/Z+H$ channel (Associated Production)

Tevatron/CDF - Associated Production

Tevatron Run II Preliminary



$q\bar{q} \rightarrow WH$ with $H \rightarrow b\bar{b}$



$W \rightarrow q\bar{q}$ 70%

- final state $q\bar{q}b\bar{b}$
- Four jet backgrounds still too large



$W \rightarrow e\nu_e$ 10%

$W \rightarrow \mu\nu_\mu$ 10%

- Final state $l\nu b\bar{b}$


- One electron or muon
- Missing transverse momentum
- Two jets
- One or two b -tags

- Easy to select in trigger and offline


$\sigma \times Br \approx 0.02$ pb



Latest Higgs Results




Tevatron Run II Preliminary, $\langle L \rangle = 5.9 \text{ fb}^{-1}$



Combined from many measurements.


10 years of data



$H \rightarrow WW \rightarrow l\nu l\nu$

1 year of data-taking

Predicted Sensitivity LHC in 2011




By end of 2011 expect to exclude Higgs in $130 < M < 460$ GeV

Compare to actual result on previous page


Is the Standard Model all there is?

- So far we have assumed a Standard Model Higgs but...
 - Does not explain Dark Matter
 - Does not unify electromagnetism weak and strong forces at high-energies (10^{16} GeV, Planck mass).
 - Do not know the Higgs potential
 - Calculations of Higgs mass using Standard Model produces a mass which is far too high (>1 TeV)
- Need models beyond the Standard Model



Supersymmetry

Every particle has a “super-partner” particle



Supersymmetric Higgs

- Need at least two Higgs doublets (H_1, H_2) to generate down- and up-type particles.
- Physical particles:
 - Radiative corrections can change masses.
 - Higgs sector now described by two free parameters (m_h and $\tan\beta=v_2/v_1$).
 - However, the exact SUSY symmetry has to be broken to reconcile the theory with experiment (i.e. the standard model and SUSY particles have different masses).
 - The minimal extension to SUSY (MSSM) has 105 parameters!
 - Have to assume a specific model e.g. mSUGRA
 - **Modifies Higgs mechanism**
 - **5 free parameters:**
 - $\tan\beta$ (as before)
 - m_0 (universal scalar mass, includes Higgs)
 - $m_{1/2}$ (gaugino mass)
 - plus two others

$$h = H_2 \cos \alpha - H_1 \sin \alpha \quad (m_h < m_Z)$$

$$H = H_2 \sin \alpha - H_1 \cos \alpha \quad (m_H > m_Z)$$

A = CP-odd Higgs

$$H^\pm = \text{charged Higgs} \quad (m_{H^\pm} = m_A^2 + m_W^2)$$

Looking for SUSY Higgs at the LHC

■ Small $\tan\beta$

- $gg \rightarrow H, A$ production is enhanced due to stronger $t\bar{t}H$ coupling.
- $H, A \rightarrow t\bar{t}$ decay gets enhanced.

■ Large $\tan\beta$

- H, A production is enhanced in $b\bar{b}$ -fusion
- $H \rightarrow \tau\tau$ has a large branching ratio

■ Medium $\tan\beta$


- Only SM-like h visible. We could see a Higgs and not realise we have seen SUSY!

■ Charged Higgs


- Clear signal for new physics (not predicted in Standard Model)

What might we see?



What we know now



What the LHC might see




What we currently see (March 2011)



Looking for other SUSY particles

- SUSY predicts that every Standard Model particle has a Super-Symmetric partner
 - Electron \leftrightarrow selectron, quark \leftrightarrow squark, $W \leftrightarrow$ wino, etc...
 - But masses not the same \rightarrow SUSY not exact symmetry
- SUSY can be a new source of CP-Violation
 - Explain matter/anti-matter asymmetry of the Universe
- A SUSY particle will quickly decay to the Lightest Supersymmetric Particle (LSP).
 - Neutral (no charge)
 - LSP is a candidate for Dark Matter
- LSP will leave detector without interacting
 - Large Missing energy, momentum (because LSP is massive)
- What is the LSP?
 - Don't really know
 - Likely to be a neutralino

What a SUSY decay looks like




Missing energy


Lots of leptons produced. Easy to see and not produced in background events

SUSY particles at the LHC

Theory predictions




Real measurements 2011



Glino excluded below 520 GeV


What LHC will see

- This assumes LHC is running at its design luminosity ($10^{33} \text{ cm}^{-2}\text{s}^{-1}$) and 7 on 7 TeV energy.
- In 2010, only running at $3 \times 10^{31} \text{ cm}^{-2}\text{s}^{-1}$ and 3.5 on 3.5 TeV energy per beam.
- So will take ~ 50 times longer than in the plot.
- However, luminosity will be increased in 2011
- On 22nd April 2011 reached $4.67 \times 10^{32} \text{ cm}^{-2}\text{s}^{-1}$




The End


ATLAS detector




ATLAS beam-pipe



ATLAS construction



ATLAS Tracker (silicon)



ATLAS toroid magnet





5th May 2011

Fergus Wilson, RAL


32

CMS detector






Inserting CMS tracker



Inserting CMS tracker



Damaged magnets 2009



5th May 2011

F QQBI.27R3 L

37